

Relating spot and flare/superflare occurrence on dwarf stars

Markus Ornik

April 1, 2025



Master's Thesis

Relating spot and flare/superflare occurrence on dwarf stars

Markus Ornik

in partial fulfillment of the requirements for the degree of
Master of Science – MSc

Supervisors

Assoz. Prof. Mag. Dr.rer.nat. Manuela Stadlober-
Temmer
Mag. Dr.rer.nat. Martin Leitzinger

April 1, 2025

Markus Ornik

Relating spot and flare/superflare occurrence on dwarf stars –

Thesis in partial fulfillment of the requirements for the degree of Master of Science – MSc;

Unsubmitted Thesis Manuscript: compiled on April 1, 2025

Institute of Physics, NAWI Graz, University of Graz.

Supervisors: Assoz. Prof. Mag. Dr.rer.nat. Manuela Stadlober-Temmer¹ and Mag. Dr.rer.nat. Martin Leitzinger^{1,2}

External Readers: Dr. Hypatia¹, Doz. Dr. Galileo Galilei², and Prof. Dr. Carl Sagan^{1,2}

¹ Institut für Physik, NAWI Graz, University of Graz, Universitätsplatz 5/II, 8010 Graz, Austria.

² Instituto de Física de Canarias, E-38200 La Laguna, Tenerife, Spain.

Abstract

Context.

Aims.

Methods.

Results.

Conclusions.

Acknowledgement

I gratefully acknowledge my supervisors Assoz. Prof. Mag. Dr.rer.nat. Manuela Stadlober-Temmer and Mag. Dr.rer.nat. Martin Leitzinger for guiding me through the process.

I thank my family and friends for supporting me.

This thesis made use of the experimental setups, laboratory and technical infrastructure, general resources and observational facilities of the Institute of Physics of the University of Graz. This thesis also made use of infrastructure, which was supported by NAWI Graz.

This research made use of pandas (team, 2020; McKinney, 2010). This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration et al., 2013; Astropy Collaboration et al., 2018). This research made use of SciPy (Virtanen et al., 2020). This research made use of matplotlib, a Python library for publication quality graphics (Hunter, 2007). This research made use of NumPy (Harris et al., 2020a; Harris et al., 2020b). This research made use of Astroquery (Ginsburg et al., 2019).

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

This paper includes data collected by the Kepler mission and obtained from the MAST data archive at the Space Telescope Science Institute (STScI). Funding for the Kepler mission is provided by the NASA Science Mission Directorate. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555.

This paper includes data collected with the TESS mission, obtained from the MAST data archive at the Space Telescope Science Institute (STScI). Funding for the TESS mission is provided by the NASA Explorer Program. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555.

I gratefully acknowledge the Austrian Science Fund (FWF): P30949-N36 (PI: xxx) for supporting this project.

Contents

Abstract	v
Acknowledgements	v
1 Introduction	1
1.1 Current knowledge	1
1.2 Spectral Types	1
1.3 Flares and Starspots	1
1.4 Space missions	1
2 GUI	3
2.1 Data download	3
2.2 Data display and manipulation	5
2.3 Data processing	10
3 Data and Methods	11
3.1 Data selection	11
3.2 Data reduction algorithms	11
3.2.1 This is a subsection	11
3.3 Example of a code snippet	11
4 Results	15
5 Discussion	17
6 Conclusion and Outlook	19
Bibliography	21
Appendix	
A Literature values for <i>Kepler</i> systems	23
A.1 Compilation of literature values	23
A.2 Tidal properties of selected systems	27
B Example for the presentation of programming code	29

B.1	System dependencies	29
B.2	Program to name new Planets	30
B.3	Example for long code listing	30
C	Appendix C	35
	Curriculum Vitae	37
	List of publications	40

Introduction

1.1 Current knowledge

1.2 Spectral Types

1.3 Flares and Starspots

1.4 Space missions

Thompson et al. (2012).

Kepler 6

$$c^2 = a^2 + b^2, \quad (1.1)$$

$$c = \sqrt{a^2 + b^2}, \quad (1.2)$$




Fig. 1.1.: Position of the 18 red giant heartbeat stars from Tables 1 and 2 in the HR diagram, where the colour shows the mass of the red giant, derived from seismology. The size of the dots represents the orbital period, ranging between 20 and 438 d. The contour surfaces reflect the density distribution of 1000 pulsating red giants. The darkest areas mark the position of the densely populated red clump. Numbers in red indicate the star count per bin, for which the contour surfaces have been drawn. Lines of equal radii in the HR diagram have been drawn for selected stellar radii between 5 and 30 $R_{\odot}$. (Figure taken from Beck et al., 2014)

GUI

In this chapter the usage of the developed GUI application is explained. It was developed to easily download new data, as well as try various algorithms for flare detection and the usage of various functions of the `lightkurve` python package (Lightkurve Collaboration et al. (2018)).

2.1 Data download

During the first startup the GUI window (e.g. in figure 2.2) is nearly completely empty. An empty local database will be created with SQLite3 (SQLite Development Team (2025)). To show lightcurves, first data needs to be downloaded. For that, the "Add" button has to be clicked. A new dialog window will show. It can be seen in figure 2.1. This GUI uses the `astroquery` (Ginsburg et al. (2019)) python package to fetch metadata and download fits files from MAST (Mikulski Archive for Space Telescopes).

The first step is to enter the identifier of the star(s). In case of multiple, they have to be separated by a semicolon. Then the missions from which the data should be fetched can be selected. The supported missions are TESS and Kepler/K2. For Kepler and K2 short and/or long cadence can be selected. The differences are described in section 1.4. After the selection has been made, the fetch button needs to be pressed. The program will then request metadata matching the star identifier(s) and selected missions from the MAST archive. The found mission data is then displayed in the preview field. It can then be downloaded by pressing the "Ok" button. The program then proceeds to download the according fits-files. The filename and TESS sector or Kepler/K2 target table ID is appended to the previously fetched metadata and handed back to the main GUI.

There a request to Simbad is made to fetch additional metadata. First the alternative identifiers are fetched from Simbad, which also has a `MAIN_ID` attribute. This is to make sure we only have one entry in the local database per star and no duplicates. Afterwards additional data is fetched from Simbad (Wenger et al. (2000)) like spectral type, radial velocity and distance. After the data is gathered, the main identifier, the alternative identifiers, spectral type, rotational velocity and distance (and their respective units) as well as the file path, source (TESS, Kepler or K2),




Fig. 2.1.: GUI window for downloading new fit files with astroquery designed using the Qt5 designer shipped with PyQt5 (Riverbank Computing Limited (2025)). The star identifier field supports one or more (separated by a semicolon) identifiers. There are checkboxes to select TESS, Kepler (long and/or short cadence) and K2 (long and/or short cadence) data. The preview field shows metadata (Identifier, TESS quarter or Kepler/K2 target table ID) to check the data which was found and which can be downloaded. The "Ok" button then proceeds to download the data and hands information back to the main GUI.

and sector/target table id are saved into a local database using SQLite3 (SQLite Development Team (2025)).

2.2 Data display and manipulation

Afterwards the list of stars seen in figure 2.2 will be refreshed. It shows the via Simbad determined main identifier alphabetically sorted.

When one of the stars in the list is selected, the program will show the additional saved informations. This includes all known IDs (in the Alt. IDs list), the spectral type, its rotational velocity and its distance. Additionally a fit type for the folded lightcurves which are described in section 3.2. The default for this is "sine". The list to the right of the information section shows the available fit files. The formatting for the list entries is "TESS - <sequence>" and "Kepler/K2 - <target table id>". The combine button allows to show multiple fit files at once if multiple are selected. Select one or combining multiple fit files, enables the plot options.

The main plot options allow the user to change the type of flux thats plotted. This is either "SAP_FLUX" (simple aperture photometry) or "PDCSAP_FLUX" (presearch data conditioning simple aperture photometry) (Lightkurve Collaboration et al. (2018)). The PDCSAP_FLUX has parts of its data removed depending on the quality flags (e.g. indicating systematic errors like an attitude tweak or a cosmic ray (Twicken et al. (2020), Thompson et al. (2016))) as well as corrected long-term brightness changes (Lightkurve Collaboration et al. (2018)). An example of a PDCSAP_FLUX is shown in figure 2.2.

The lightkurve python package also supports normalizing (to unscaled, percent, parts per thousands (ppt) or parts per million (ppm)) which can be seen in figure 2.4. Flattened lightcurves (see figure 2.5 as an example for *BD – 08 995*) is used to identify flares. The "remove outliers", "remove nans/infs" and "bin" features are exposed in the GUI, but are not actively used in this thesis.

The checkboxes to flatten the lightcurve (figure 2.5), fold the lighhtcurve (figure 2.6) and showing the periodogram (figures 2.7 and 2.8) are exclusive of each other. Only one can be used at a time.




Fig. 2.2.: Main GUI window. This window was designed with the Qt5 designer shipped with PyQz5 (Riverbank Computing Limited (2025)). It shows a list of all available stars with their main identifier (taken from SIMBAD (Wenger et al. (2000))) sorted alphabetically. The "Star infos" box contains additional information on the star gathered from SIMBAD. There is a list of alternative IDs, the spectral type, the rotational velocity, and the distance. It also shows the preferred fit type for folding lightcurves. On the right of the information box is a list of all available lightcurves, with an option to combine multiple to plot at once. To the right of the list of available lightcurves are multiple different plot options, which utilize the different capabilities of the lightcurve python package (Lightkurve Collaboration et al. (2018)). The values for "Period" and "Epoch Time" are automatically calculated and used as default for the selected lightcurve. In this image the TESS Sector 32 lightcurve of *BD* – 08 995 is shown. The red crosses show the peaks of the found flares, while the red overall over the lightcurve indicates the duration of the flare.




Fig. 2.3.: Main GUI window. Same as in figure 2.2, with the difference that the data quality flag is also shown.




Fig. 2.4.: Main GUI window. Same as in figure 2.2, with the difference that the lightcurve was normalized. It also shows all possible normalize options the lightcurve python package (Lightcurve Collaboration et al. (2018)) supports.




Fig. 2.5.: Main GUI window. Showing the same lightcurve as figure 2.2, with the difference that it is now flattened. The green line shows a fit of the flare. This is described in chapter 3.2.




Fig. 2.6.: Main GUI window. Showing the same lightcurve as figure 2.2, with the difference that it is now folded with an period of 2.697 days. Additionally vertical lines are shown which indicate the region around the minima and maxima of the folded lightcurve respectively.




Fig. 2.7.: Main GUI window. Showing the periodogram of the lightcurve in figure 2.2 using the lombscargle algorithm. Furthermore it shows the four highest peaks (which are atleast 100 datapoints separated). The method is explained in chapter 3.2.




Fig. 2.8.: Main GUI window. Showing the periodogram of the lightcurve in figure 2.2 using the box least square algorithm. Similarly to figure 2.7 the four highest peaks are marked with a red "x".

2.3 Data processing

Pressing the button "New FC" will start the processing of all fits files in the current loaded database. It will sequentially go through all files and apply the flare detection algorithms to the lightcurve, as well as generate the periodogram and folded lightcurves. This is done for both the SAP_FLUX as well as the PDCSAP_FLUX. Additionally plots for each individual lightcurve and processing step are generated. The included data includes:

- Flares (peak, timestamp, datapoint index in the lightcurve, TESS/Kepler data quality flags)
- Number of flares
- Fits for the flares (fit, timestamps, indices in the lightcurve, TESS/Kepler data quality flags)
- Uninterrupted timespans in the data
- Total length of timespans in seconds
- Folded lightcurve phase range
- Folded lightcurve flare positions (phase, peak)
- Period which was used to fold the lightcurve
- Additional boundaries around minima and maxima

After this process is done, a new window will open. It allows to show multiple different statistics like flares per file, flares per star (total, or normalized to per 7 day period) or mean periods for each star. This window can be seen in figure ??.

Table 3.1.: Training sample of false-positive planet candidates for the clustering algorithm.

Star's identifier	T_{eff} [K]	Lum [$L/L_{\odot}$]	Luminosity cluster	Reference
π Her	4170	1320	High	Hatzes and Cochran (1999)
μ UMa	3899	1150	High	Lee et al. (2016)
HD 18438	3871	830	High	Delgado Mena et al. (2018)
γ Dra (Eltanin)	3990	510	High	Hatzes et al. (2018)
α Tau (Aldebaran)	4055	440	High	Hatzes et al. (2015)
θ^1 Tau	5000	70	Low	Beck et al. (2015)
NGC2423 No.3	4592	65	Low	Delgado Mena et al. (2018)
γ Psc	4909	60	Low	Beck et al. (2015)
β Gem (Pollux)	4865	40	Low	Delgado Mena et al. (2018)
IC4651 No.9122	4720	35	Low	Delgado Mena et al. (2018)
γ Cep A	4900	10	Low	Hatzes and Cochran (1999)

Notes. The star's identifier is given. For the most prominent stars, the name of the star is amended in brackets. The next columns report on the effective temperature and the luminosity of the star in units of the solar luminosity $L_{\odot}$. The typical uncertainty of the reported temperature and luminosity are 150K and $\sim 10\%$, respectively. The top and bottom panel report classification of the clustering. The cluster name refers to the position in the red-giant phase. The literature references are provided in the last column. (Table taken from Elisabeth Höldrich bachelor thesis, 2020)

3

Data and Methods

In this chapter the selection criteria for the data is explained in section 3.1. Furthermore a detailed description of the algorithm is given in section 3.2.

3.1 Data selection

3.2 Data reduction algorithms

3.2.1 This is a subsection

3.3 Example of a code snippet

Listing 3.1: Example for setting the tidal mechanisms in the *star_star* inlist.

```
1 &binary_controls
2     do_tidal_circ = .true.
3     circ_type_1 = 'Hut_conv'
4     circ_type_2 = 'Hut_conv'
5
6     do_tidal_sync = .false.
7     sync_type_1 = 'Hut_conv'
8     sync_type_2 = 'Hut_conv'
```

Notes. A part of *&binary_controls* environment in the *star_inlist* is shown. The true/false-statements in line 2 and 6 determine if the models defined below takes into account the defined mechanism. The keywords for defining the formalistic description for circularization and synchronization mechanism are defined in lines 2-4 and 6-8, respectively. The number at the end of the keyword indicates which formalism is set for the primary (1) and secondary (2) stellar component.




Fig. 3.1.: Disentangled spectra of KIC 9163796. The spectra of the primary in blue and secondary in red are shown in the left and right spectrum, respectively. Top panel: the region around the $H\beta$ line, while bottom panel: magnesium triplet at 518 nm, normalised to the continuum flux of the composite spectrum. The synthetic model of the best fit of the determination of fundamental parameters and metallicity is shown as dashed grey line is shown depicted in the region of the Mg triplet. (Graphic taken from Beck et al., 2018)

Table 3.2.: Seismic and fundamental parameters for 14 oscillating red giant heartbeat stars, ordered by descending orbital period.

Star KIC	$\nu_{\max}$ [μHz]	$\Delta\nu$ [μHz]	$\Delta\Pi_1$ [sec]	$\delta f_{\max}$ [nHz]	Evol. Phase	R [$R_{\odot}$]	M [$M_{\odot}$]	$\log g$ [dex]	L [$L_{\odot}$]	T_{eff} [K]	P_{orbit} [d]	A [ppt]	$ \Delta\text{RV} $ [km s^{-1}]
9151763	13.8 $\pm$ 0.2	1.98 $\pm$ 0.01	—	—	RGB?	17.6 $\pm$ 0.4	1.19 $\pm$ 0.08	2.01	96 $\pm$ 16	4290	437.5	+7.1	32.2
7431665	54.0 $\pm$ 0.7	5.46 $\pm$ 0.02	$\sim$ 67	—	RGB	9.4 $\pm$ 0.1	1.39 $\pm$ 0.05	2.62	35 $\pm$ 2	4580	281.4	-3.0	[37.8]
5039392	6.2 $\pm$ 0.1	1.13 $\pm$ 0.01	—	—	RGB	24.0 $\pm$ 0.7	0.98 $\pm$ 0.07	1.67	157 $\pm$ 24	4110	236.7	-6.0	42.3
9540226*	27.4 $\pm$ 0.4	3.18 $\pm$ 0.01	—	—	RGB	14.1 $\pm$ 0.3	1.6 $\pm$ 0.1	2.37	81 $\pm$ 13	4600	175.4	-7	45.3
8210370	44.1 $\pm$ 0.8	4.69 $\pm$ 0.02	—	—	RGB?	10.5 $\pm$ 0.2	1.40 $\pm$ 0.08	2.54	44 $\pm$ 4	4585	153.5	-5.3	22.1
11044668	50.2 $\pm$ 0.2	5.65 $\pm$ 0.01	$\sim$ 60	83(?)	RGB	8.18 $\pm$ 0.09	0.99 $\pm$ 0.03	2.59	26 $\pm$ 3	4565	139.5	-3.8	[43.0]
10614012*	70.2 $\pm$ 0.9	6.54 $\pm$ 0.02	—	—	RGB	8.6 $\pm$ 0.2	1.49 $\pm$ 0.08	2.74	33 $\pm$ 4	4715	132.1	-4.7	49.3
9163796	153.2 $\pm$ 0.7	13.53 $\pm$ 0.04	—	—	RGB	4.46 $\pm$ 0.03	0.89 $\pm$ 0.01	3.09	12 $\pm$ 1	4820	121.3	$\pm$ 0.5	70.1
2444348	30.5 $\pm$ 0.3	3.26 $\pm$ 0.01	—	—	RGB	14.9 $\pm$ 0.3	1.94 $\pm$ 0.11	2.38	86 $\pm$ 14	4565	103.5	-1.7	7.7
5006817	145.9 $\pm$ 0.5	11.64 $\pm$ 0.01	78	450	RGB	5.84 $\pm$ 0.09	1.49 $\pm$ 0.06	3.08	19 $\pm$ 3	5000	94.8	-1.7	23.5
8803882	347 $\pm$ 3	22.6 $\pm$ 0.4	—	500(?)	RGB	3.68 $\pm$ 0.1	1.4 $\pm$ 0.1	3.45	8 $\pm$ 1	5043	89.7	+0.5	[1.9]
8144355	179 $\pm$ 2	13.95 $\pm$ 0.04	$\sim$ 78	210(?)	RGB	4.90 $\pm$ 0.09	1.26 $\pm$ 0.08	3.16	12 $\pm$ 2	4875	80.6	+2.1	18.9
9408183	164.8 $\pm$ 0.2	13.29 $\pm$ 0.02	$\sim$ 93	450	RGB	5.02 $\pm$ 0.07	1.23 $\pm$ 0.05	3.12	13 $\pm$ 1	4900	49.7	+1.5	64.4
2720096	110.1 $\pm$ 0.7	9.17 $\pm$ 0.01	—	—	RGB	6.98 $\pm$ 0.08	1.54 $\pm$ 0.06	2.95	23 $\pm$ 2	4812	26.7	+1.0	4.0
8095275	69.3 $\pm$ 0.3	6.81 $\pm$ 0.01	—	—	RGB	7.78 $\pm$ 0.08	1.21 $\pm$ 0.05	2.74	25 $\pm$ 3	4622	23.0	-6.0	20.6

Notes. The star's identifier in the *Kepler* Input Catalogue (KIC) is given. Eclipsing systems are marked with an asterisk. The columns $\nu_{\max}$ and $\Delta\nu$ report the frequency of the oscillation power excess and the large frequency separation between radial modes for a given star. $\Delta\Pi_1$ quantifies the true period spacing of dipole modes. The maximum value of the detected rotational splitting δf is listed. The evolutionary phase RGB describes H-shell burning red giant. Ambiguous values are marked with '?'. The columns R , M , L , and $\log g$ report the stellar radius, mass, luminosity, effective temperature and surface gravity from scaling relations, respectively. T_{eff} was adopted from the KIC. The uncertainties of $\log g$ are on the order of 0.01 dex and for the temperature typically smaller than 150 K. P_{orbit} gives the orbital period from photometry. The column A lists the maximum amplitude of the heartbeat in a rebinned phase diagram. The error estimate for P_{orbit} and A from the PDM is not reliable due to the remaining contamination of the solar-like oscillations and therefore not given. $|\Delta\text{RV}|$ reports the maximum difference in radial velocity. Squared brackets mark systems for which the orbital parameters could not yet be determined from radial velocities. (Table taken from Beck et al., 2014)

Results

4

Discussion

5

Conclusion and Outlook

Bibliography

- Astropy Collaboration, A. M. Price-Whelan, B. M. SipHocz, et al. (Sept. 2018). „The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package“. In: *aj* 156.3, 123, p. 123. DOI: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f). arXiv: [1801.02634](https://arxiv.org/abs/1801.02634) [astro-ph.IM] (cit. on p. vii).
- Astropy Collaboration, T. P. Robitaille, E. J. Tollerud, et al. (Oct. 2013). „Astropy: A community Python package for astronomy“. In: *Astronomy and Astrophysics* 558, A33, A33. DOI: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068). arXiv: [1307.6212](https://arxiv.org/abs/1307.6212) [astro-ph.IM] (cit. on p. vii).
- Beck, P. G., T. Kallinger, K. Pavlovski, et al. (Apr. 2018). „Seismic probing of the first dredge-up event through the eccentric red-giant and red-giant spectroscopic binary KIC 9163796. How different are red-giant stars with a mass ratio of 1.015?“ In: *Astronomy and Astrophysics* 612, A22, A22. DOI: [10.1051/0004-6361/201731269](https://doi.org/10.1051/0004-6361/201731269). arXiv: [1712.05208](https://arxiv.org/abs/1712.05208) [astro-ph.SR] (cit. on pp. 12, 23).
- Beck, P. G., E. Kambe, M. Hillen, et al. (Jan. 2015). „Detection of solar-like oscillations in the bright red giant stars γ Piscium and θ^1 Tauri from a 190-day high-precision spectroscopic multi-site campaign“. In: *Astronomy and Astrophysics* 573, A138, A138. DOI: [10.1051/0004-6361/201323019](https://doi.org/10.1051/0004-6361/201323019). arXiv: [1407.6352](https://arxiv.org/abs/1407.6352) [astro-ph.SR] (cit. on p. 11).
- Beck, P.G., K. Hambleton, J. Vos, et al. (2014). „Pulsating red giant stars in eccentric binary systems discovered from Kepler space-based photometry-A sample study and the analysis of KIC 5006817“. In: *Astronomy and Astrophysics* 564, A36. DOI: [10.1051/0004-6361/201322477](https://doi.org/10.1051/0004-6361/201322477) (cit. on pp. 2, 13).
- Delgado Mena, E., C. Lovis, N. C. Santos, et al. (Nov. 2018). „Planets around evolved intermediate-mass stars. II. Are there really planets around IC 4651 No. 9122, NGC 2423 No. 3, and NGC 4349 No. 127?“ In: *Astronomy and Astrophysics* 619, A2, A2. DOI: [10.1051/0004-6361/201833152](https://doi.org/10.1051/0004-6361/201833152). arXiv: [1807.09608](https://arxiv.org/abs/1807.09608) [astro-ph.EP] (cit. on p. 11).
- Ginsburg, A., B. M. Sipőcz, C. E. Brasseur, et al. (Mar. 2019). „astroquery: An Astronomical Web-querying Package in Python“. In: *The Astronomical Journal* 157, 98, p. 98. DOI: [10.3847/1538-3881/aafc33](https://doi.org/10.3847/1538-3881/aafc33). arXiv: [1901.04520](https://arxiv.org/abs/1901.04520) [astro-ph.IM] (cit. on pp. vii, 3).
- Harris, Charles R., K. Jarrod Millman, Stéfan J. van der Walt, et al. (Sept. 2020a). „Array programming with NumPy“. In: *Nature* 585.7825, pp. 357–362. DOI: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2) (cit. on p. vii).
- Harris, Charles R., K. Jarrod Millman, Stéfan J. van der Walt, et al. (Sept. 2020b). „Array programming with NumPy“. In: *Nature* 585.7825, pp. 357–362. DOI: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2) (cit. on p. vii).

- Hatzes, A. P., W. D. Cochran, M. Endl, et al. (Aug. 2015). „Long-lived, long-period radial velocity variations in Aldebaran: A planetary companion and stellar activity“. In: *Astronomy and Astrophysics* 580, A31, A31. DOI: [10.1051/0004-6361/201425519](https://doi.org/10.1051/0004-6361/201425519). arXiv: [1505.03454](https://arxiv.org/abs/1505.03454) [astro-ph.SR] (cit. on p. 11).
- Hatzes, A. P., M. Endl, W. D. Cochran, et al. (Mar. 2018). „The Radial Velocity Variability of the K-giant γ Draconis: Stellar Variability Masquerading as a Planet“. In: *The Astronomical Journal* 155.3, 120, p. 120. DOI: [10.3847/1538-3881/aaa8e1](https://doi.org/10.3847/1538-3881/aaa8e1). arXiv: [1801.05239](https://arxiv.org/abs/1801.05239) [astro-ph.SR] (cit. on p. 11).
- Hatzes, Artie P. and William D. Cochran (Mar. 1999). „Long-period, low-amplitude radial velocity variations in the K giant star π Herculis: rotation, substellar companion or non-radial pulsations?“ In: *Monthly Notices of the RAS* 304.1, pp. 109–118. DOI: [10.1046/j.1365-8711.1999.02288.x](https://doi.org/10.1046/j.1365-8711.1999.02288.x) (cit. on p. 11).
- Hunter, J. D. (2007). „Matplotlib: A 2D Graphics Environment“. In: *Computing in Science Engineering* 9.3, pp. 90–95 (cit. on p. vii).
- Lee, Byeong-Cheol, Inwoo Han, Myeong-Gu Park, et al. (Apr. 2016). „Long-period Variations in the Radial Velocity of Spectroscopic Binary M Giant μ Ursae Majoris“. In: *The Astronomical Journal* 151.4, 106, p. 106. DOI: [10.3847/0004-6256/151/4/106](https://doi.org/10.3847/0004-6256/151/4/106). arXiv: [1602.07011](https://arxiv.org/abs/1602.07011) [astro-ph.SR] (cit. on p. 11).
- Lightcurve Collaboration, J. V. d. M. Cardoso, C. Hedges, et al. (Dec. 2018). *Lightcurve: Kepler and TESS time series analysis in Python*. Astrophysics Source Code Library. ascl: [1812.013](https://ascl.net/1812.013) (cit. on pp. 3, 5–7).
- McKinney, Wes (2010). „Data Structures for Statistical Computing in Python“. In: *Proceedings of the 9th Python in Science Conference*. Ed. by Stéfan van der Walt and Jarrod Millman, pp. 56–61. DOI: [10.25080/Majora-92bf1922-00a](https://doi.org/10.25080/Majora-92bf1922-00a) (cit. on p. vii).
- Riverbank Computing Limited (2025). *PyQt5* (cit. on pp. 4, 6).
- SQLite Development Team (2025). *SQLite* (cit. on pp. 3, 4).
- team, The pandas development (Feb. 2020). *pandas-dev/pandas: Pandas*. Version latest. DOI: [10.5281/zenodo.3509134](https://doi.org/10.5281/zenodo.3509134) (cit. on p. vii).
- Thompson, Susan E., Mark Everett, Fergal Mullally, et al. (2012). „A Class of Eccentric Binaries with Dynamic Tidal Distortions Discovered with Kepler“. In: *The Astrophysical Journal* 753.1, p. 86 (cit. on p. 1).
- Thompson, Susan E., Dorothy Fraquelli, Jeffrey E. Van Cleve, and Douglas A. Caldwell (May 2016). „Kepler Archive Manual“. In: (cit. on p. 5).
- Twicken, Joseph D., Douglas A. Caldwell, Jon M. Jenkins, et al. (Sept. 2020). „TESS Science Data Products Description Document“. In: (cit. on p. 5).
- Virtanen, Pauli, Ralf Gommers, Travis E. Oliphant, et al. (2020). „SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python“. In: *Nature Methods* 17, pp. 261–272. DOI: <https://doi.org/10.1038/s41592-019-0686-2> (cit. on p. vii).
- Wenger, M., F. Ochsenbein, D. Egret, et al. (Apr. 2000). „The SIMBAD astronomical database. The CDS reference database for astronomical objects“. In: *Astronomy and Astrophysics Supplement Series* 143, pp. 9–22. DOI: [10.1051/aas:2000332](https://doi.org/10.1051/aas:2000332). arXiv: [astro-ph/0002110](https://arxiv.org/abs/astro-ph/0002110) [astro-ph] (cit. on pp. 3, 6).

Literature values for *Kepler* systems

[This appendix has been published as Appendix C of Beck et al. (2018)].

The parameters for eclipsing binaries are taken from the dynamical solution of G16. For the list of heartbeat stars (B14), radius and mass were inferred from seismic scaling relations and corrected for the systematic mass overestimate of 15% reported by G16. Surface rotation periods were adopted from G14, B14 and B18. For four stars of B14, KIC 7431665, KIC 11044668, KIC 8803882, and KIC 7799540, no orbital parameters have yet been published.

A.1 Compilation of literature values

We refer to the cited literature for details on the applied methodology. Table A.1 contains the following parameters,

- *KIC* specifies the target identification number in the *Kepler* Input Catalog.
- *Type* indicates if a binary system is an eclipsing binary (EB), a heartbeat system (HB), or an eclipsing heartbeat system (eHB). 'NO' indicates that the star belongs to the four non-oscillating stars of G14/G16.
- P_{orbit} is the measured orbital period.
- e is the orbital eccentricity.
- ν_{max} is the peak frequency of the excess of oscillation power.
- $R/R_{\odot}$ is the stellar radius in solar units. Values from B14 are seismically inferred and corrected for the 5% overestimate of seismic radius. Values from G16 originate from a dynamical solution.

- $M/M_{\odot}$ is the stellar mass in solar units. Values from B14 are seismically inferred and corrected for the 15% overestimate of seismic mass. Values from G16 originate from a dynamical solution.
- $q = M_2/M_1$ is the mass ratio between the two stellar components in the system. '?' indicates if q has not been determined for a given system.
- T_{eff} is the effective temperature.
- P_{rot} specifies the time scale of the flux modulation, identified as the surface rotation period. The sources of the values are G14 and B18. We round all period values to the next full day.
- The dimensionless number ε_r is proportional to the inverse of the time scale of tidal circularisation (see Eq. XX). If no value of the mass ratio is specified, ε_r is calculated for $q = 0.5$.
- REF: the last column is specifying the literature references. If several papers are reporting on a given system, values of the most recent paper are cited. Previous references are given in brackets.

Table A.1.: Literature Parameters of red-giant binaries in the *Kepler* sample.

KIC	Type	P _{orb} [days]	<i>e</i> []	$\nu_{\max}$ [μ Hz]	R/R _☉ []	M/M _☉ []	<i>q</i> []	T [K]	P _{rot} [days]	ε_R []	REF
2444348	HB	103.50 ± 0.01	0.48 ± 0.01	30.5 ± 0.3	14.2 ± 0.3	1.6 ± 0.1		? 4565	-	-0.53	B14
2697935	eHB	21.50 ± 0.02	0.41 ± 0.02	~ 405.6	~ 3.1	~ 1.2		? 4883	-	-0.73	B14
2720096	HB	26.70 ± 0.01	0.49 ± 0.01	110.1 ± 0.7	6.6 ± 0.1	1.3 ± 0.1		? 4812	-	0.83	B14
3955867	EB, NO	33.65685 ± 0.00007	0.019 ± 0.002	–	7.9 ± 0.1	1.10 ± 0.06	0.84 ± 0.05	4884	33	1.14	G16 (G14)
4569590	EB, NO	41.3710 ± 0.0001	0.004 ± 0.001	–	14.1 ± 0.2	1.6 ± 0.10	0.66 ± 0.05	4706	41	1.67	G16 (G14)
4663623	EB	358.09 ± 0.0003	0.43 ± 0.01	54.1 ± 0.2	9.7 ± 0.2	1.36 ± 0.09	0.99 ± 0.08	4812	-	-4.08	G16 (G14)
5006817	HB	94.812 ± 0.002	0.71 ± 0.01	145.9 ± 0.5	5.5 ± 0.1	1.3 ± 0.1	0.199 ± 0.001	5000	-	-2.80	B14
5039392	HB	236.70 ± 0.02	0.44 ± 0.01	6.2 ± 0.1	22.8 ± 0.7	0.8 ± 0.1		? 4110	-	-0.01	B14
5179609	EB	43.93108 ± 0.000002	0.150 ± 0.001	322 ± 1.0	3.50 ± 0.03	1.18 ± 0.03	0.51 ± 0.02	5003	182	-1.96	G16 (G14)
5308778	EB	40.5661 ± 0.0003	0.006 ± 0.005	49 ± 1.1	10.1 ± 0.3	1.5 ± 0.1	0.43 ± 0.03	4900	39	0.80	G16 (G14)
5786154	EB	197.918 ± 0.0004	0.3764 ± 0.0009	29.8 ± 0.2	11.4 ± 0.2	1.06 ± 0.06	0.96 ± 0.07	4747	-	-1.85	G16 (G14)
7037405	EB	207.1083 ± 0.0007	0.238 ± 0.004	21.8 ± 0.1	14.1 ± 0.2	1.25 ± 0.04	0.91 ± 0.03	4516	-	-1.62	G16 (G14)
7377422	EB	107.6213 ± 0.0004	0.4377 ± 0.0005	40 ± 2.1	9.5 ± 0.2	1.05 ± 0.08	0.81 ± 0.07	4938	55	-0.96	G16 (G14)
7943602	EB, NO	14.69199 ± 0.00004	0.001 ± 0.003	–	6.6 ± 0.2	1.0 ± 0.10	0.78 ± 0.09	5096	15	2.70	G16 (G14)
8054233	EB	1058.16 ± 0.02	0.2718 ± 0.0004	46.5 ± 0.3	10.7 ± 0.1	1.60 ± 0.06	0.69 ± 0.04	4971	-	-6.61	G16 (G14)
8095275	HB	23.00 ± 0.01	0.32 ± 0.01	69.3 ± 0.3	7.4 ± 0.1	1.0 ± 0.1		? 4622	-	1.86	B14
8144355	HB	80.55 ± 0.01	0.76 ± 0.01	179.0 ± 2	4.7 ± 0.1	1.1 ± 0.1		? 4875	-	-2.41	B14
8210370	HB	153.50 ± 0.01	0.70 ± 0.01	44.1 ± 0.8	10.0 ± 0.2	1.2 ± 0.1		? 4585	-	-1.92	B14
8410637	EB	408.3 ± 0.5	0.689 ± 0.001	46.0 ± 0.2	10.7 ± 0.1	1.56 ± 0.3	0.85 ± 0.16	4800	-	-4.34	F13
8430105	EB	63.32713 ± 0.00003	0.2564 ± 0.0002	76.7 ± 0.6	7.65 ± 0.05	1.31 ± 0.02	0.63 ± 0.01	5042	122	-0.73	G16 (G14)

Table A.1.: Literature Parameters of red-giant binaries in the *Kepler* sample. (continued)

KIC	Type	P _{orb} [days]	<i>e</i> []	$\nu_{\max}$ [μ Hz]	R/R _☉ []	M/M _☉ []	<i>q</i> []	T [K]	P _{rot} [days]	ε_r []	REF
8702921	EB	19.38446 ± 0.00002	0.0964 ± 0.0008	195.6 ± 0.5	5.32 ± 0.05	1.67 ± 0.05	0.16 ± 0.01	5058	98	0.26	G16 (G14)
8912308	HB	20.17 ± 0.01	0.23 ± 0.01	~ 350.0	~ 4.0	~ 1.7		? 4872	-	-0.39	B14
9151763	HB	437.51 ± 0.03	0.73 ± 0.01	13.8 ± 0.2	16.7 ± 0.4	1.0 ± 0.1		? 4290	-	-2.62	B14
9163796	HB	121.30 ± 0.01	0.69 ± 0.002	165.3 ± 1.3	5.1 ± 0.1	1.2 ± 0.1	0.985 ± 0.005	4960	130	-3.17	B18 (B14)
9246715	EB	171.27688 ± 0.00001	0.3559 ± 0.0003	106.4 ± 0.8	8.30 ± 0.04	2.149 ± 0.007	0.990 ± 0.005	5030	93	-3.54	R16 (G14)
9291629	EB, NO	20.68643 ± 0.00004	0.007 ± 0.002	–	7.99 ± 0.05	1.14 ± 0.03	0.96 ± 0.03	4713	21	2.26	G16 (G14)
9408183	HB	49.70 ± 0.01	0.42 ± 0.01	164.8 ± 0.2	4.8 ± 0.1	1.0 ± 0.1		? 4900	-	-1.18	B14
9540226	eHB	175.4439 ± 0.0006	0.3880 ± 0.0002	27.1 ± 0.2	12.8 ± 0.1	1.33 ± 0.05	0.74 ± 0.04	4692	-	-1.64	G16 (B14, G14)
9970396	EB	235.2985 ± 0.0002	0.194 ± 0.007	63.7 ± 0.2	8.0 ± 0.2	1.14 ± 0.03	0.89 ± 0.03	4916	-	-3.38	G16 (G14)
10001167	EB	120.3903 ± 0.0005	0.159 ± 0.003	19.9 ± 0.1	12.7 ± 0.3	0.81 ± 0.05	0.98 ± 0.07	4700	-	0.03	G16 (G14)
10614012	eHB	132.13 ± 0.01	0.71 ± 0.01	70.2 ± 0.9	8.2 ± 0.2	1.3 ± 0.1		? 4715	-	-2.23	B14

A.2 Tidal properties of selected systems

Table A.2 lists the following parameters for the systems' red-giant primary with known P_{rot} ,

- δ_{10} is the 10th percentile of the logarithm of the ratio between the dissipation of the equilibrium and the dynamical tide, $\delta = \log(\mathcal{D}_{\text{eq}} / \langle \mathcal{D} \rangle_{\omega})$.
- $\tau_{\text{conv},10}$ is the corresponding convective turnover timescale computed in the middle of the convective zone.
- $P_{\text{tide}} > \tau_{\text{conv},10}$ indicates if the tidal period is longer than the convective turnover timescale (Yes / No).

Table A.2.: Evolution of δ_{10} for the systems with know rotation period.

KIC	P_{tide} [days]	$\tau_{\text{conv},10}$ [days]	$P_{\text{tide}} > \tau_{\text{conv},10}$	δ_{10}
7943602	357	20	Y	2.3-3.4
7377422	56	20	Y	3.7-4.8
3955867	845	20-23	Y	2.6-3.7
9291629	692	20-23	Y	2.3-3.4
5179609	28	23	Y	4.7-5.8
9163796	906	23	Y	3.8-4.9
8430105	65	23-26	Y	4.4-5.5
5308778	505	12	Y	3.0-4.1
4569590	2285	12	Y	2.4-3.5
8702921	12	12	N	4.2-5.3
9246715	101	14	Y	4.1-5.2

Example for the presentation of programming code

B.1 System dependencies

The presented code is written for Python 3.7. Table B.1 details the used standard and specialized python packages and modules and is structured as follows.

- The package and module names are listed in the first column. Specialized modules are preceded with the abbreviation of the parent package, whose abbreviated package name is indicated in brackets.
- The version number of the used library is indicate in the second column.
- A brief module description is provided in the third column.

The top panel reports the standard python packages, contained in the standard python distribution. The bottom panel lists specialized packages, which require individual download. (Modified table based on Table A.1 from Rafael Goldgruber's Bachelor thesis).

Table B.1.: Required Python modules for the presented code.

Package / Mod- ule	Version	Modul Description
math	—	Access mathematical functions;
os	—	Miscellaneous operating system interfaces
glob	—	Creates iterable lists from folder content
matplotlib	3.3.1	Core package for scientific computation & plotting
numpy	1.19.1	Core package for numerical computing
astropy (ap)	4.0.1	Community Python Library for Astronomy
ap.astroquery	0.4	Querying astronomical web databases
ap.vizier	0.4	Importing online data and catalogues, published along with papers
ap.io.pyfits		Reading and operating with FITS files
mesareader	—	Reading MESA history and profile output files

B.2 Program to name new Planets

Provide a short description of the following script and give a brief description of the key settings or input format. The example below shows how to insert the code of a whole program from its source file.

Listing B.1: HelloPlanet.py: Sophisticated code to name new worlds.

```
1 import sys , os , math
2 import numpy as np
3 import pylab as pl
4
5 # here we have a comment on the scripting
6
7 for iii in np.arange(0,42):
8     print( 'Hello_new_planet_number_{ }'.format( iii +1))
```

Table B.2.: Option menu for helloWorld.py (Listing B.1),

Type	Argument	Functionality
InCode	if xY==True	Description of what will happen if this option is used.
CmdL	-n	Description of what will happen if this option is used.
ReqIn	parameter name	Description of what will happen if this option is used.

Notes. Usage options: *InCode* curial options to be set inside the code before excecution of the program, *CmdL* options set in the command line, *ReqIn* requested user input during the execution of the program;

B.3 Example for long code listing

This section presents styles to present longer code snippets. To save printing space and cost, these are typset in a two-column format.

The combined presentation of Listing B.2 and Listing B.3 show how two snippets can be shown. Listing B.4 depicts how long code snippets or full programs can be included as multi-column and multi-page element.

Also for these listings a preamble stating the system requirements is needed.

Listing B.2: inlist1

```

1 &star_job
2     show_log_description_at_start = .true.
3     pgstar_flag = .true.
4 / ! end of star_job namelist
5
6 &controls
7     extra_terminal_output_file = 'log1'
8     log_directory = 'LOGS1'
9 / ! end of controls namelist
10
11 &pgstar
12     History_Panels1_win_flag = .true.
13     History_Panels1_win_width = 5
14     History_Panels1_win_aspect_ratio = 1.0
15
16     History_Panels1_title = 'Orbital_evolution'
17     History_Panels1_num_panels = 2
18
19     History_Panels1_yaxis_name(1) = 'period_days'
20     History_Panels1_other_yaxis_name(1) =
21     'lg_mstar_dot_1'
22     History_Panels1_yaxis_name(2) = 'Jdot'
23     History_Panels1_other_yaxis_name(2) =
24     'binary_separation'
25 / ! end of pgstar namelist

```

Listing B.3: inlist2

```

1 &star_job
2     show_log_description_at_start = .false.
3
4 / ! end of star_job namelist
5
6 &controls
7     extra_terminal_output_file = 'log2'
8     log_directory = 'LOGS2'
9 / ! end of controls namelist
10
11 &pgstar
12
13
14
15
16
17
18
19
20
21
22
23
24
25 / ! end of pgstar namelist

```

Listing B.4: Example for a multi-column and multi-page code listing.

```

1  ! inlist to evolve a 15 solar mass star
2
3  ! For the sake of future readers of this file ,
4  ! ONLY include the controls you are actually using.
5  ! DO NOT include all of the other controls that
6  ! simply have their default values.
7
8  &star_job
9
10 ! begin with a pre-main sequence model
11   create_pre_main_sequence_model = .true.
12
13 ! save a model at the end of the run
14   save_model_when_terminate = .false.
15   save_model_filename = '15M_at_TAMS.mod'
16
17 ! display on-screen plots
18   pgstar_flag = .true.
19
20 / !end of star_job namelist
21
22
23 &controls
24
25 ! starting specifications
26   initial_mass = 15 ! in Msun units
27
28 ! options for energy conservation
29   use_dedt_form_of_energy_eqn = .true.
30   use_gold_tolerances = .true.
31
32 ! stop when the star nears ZAMS (Lnuc/L > 0.99)
33   Lnuc_div_L_zams_limit = 0.99d0
34   stop_near_zams = .true.
35
36 ! stop when the center mass fraction
37 of h1 drops below this limit
38   xa_central_lower_limit_species(1) = 'h1'
39   xa_central_lower_limit(1) = 1d-3
40
41 / ! end of controls namelist
42
43 &pgstar
44
45 ! MESA uses PGPLOT for live plotting and
46 ! gives the user a tremendous amount of control
47 ! of the presentation of the information.
48

```

```
49 ! show HR diagram
50 ! this plots the history of L,Teff over many
51 ! timesteps
52   HR_win_flag = .true.
53
54 ! set static plot bounds
55   HR_logT_min = 3.5
56   HR_logT_max = 4.6
57   HR_logL_min = 2.0
58   HR_logL_max = 6.0
59
60 ! set window size (aspect_ratio = height/width)
61   HR_win_width = 6
62   HR_win_aspect_ratio = 1.0
63
64
```

```
65 ! show temperature/density profile
66 ! this plots the internal structure at single
67 ! timestep
68   TRho_Profile_win_flag = .true.
69
70 ! add legend explaining colors
71   show_TRho_Profile_legend = .true.
72
73 ! display numerical info about the star
74   show_TRho_Profile_text_info = .true.
75
76 ! set window size (aspect_ratio = height/width)
77   TRho_Profile_win_width = 8
78   TRho_Profile_win_aspect_ratio = 0.75
79
80 / ! end of pgstar namelist
```

Appendix C

C

Curriculum Vitae

Mag. Nikolaus Copernicus, PhD

Born	February 19, 1473
Citizenship	Poland

Research positions and academic education

Current academic position

Affiliation & academic address	Institute for Physics, Karl Franzens University Graz, Universitätsplatz 5/II, NAWI Graz, 8010 Graz, Austria
Contact	nicolaus.copernicus@uni-graz.at

Research Positions

since March 2020	Doctoral researcher at the Department for Astronomy, Inst. of Physics, NAWI Graz, University of Graz, Austria
2019, March – 2020, March	Pre-doctoral researcher at the Department for Astronomy, Inst. of Physics, NAWI Graz, University of Graz, Austria

Academic Education

2017, February – 2019, March	University of Graz, Austria: Master of Natural Sciences in Physics (Graduation: March 32, 2019), Title of Thesis
2014, October – 2017, February	University of Vienna, Austria : Bachelor of Natural Sciences in Astronomy (Graduation: February, 30, 2017)

Academic Recognition

Invited talks at conferences and seminars

- **Title of invited contribution**, title of conference or occasion, length of contribution, (date, hosting institution)

Grants & Awards

- **Name of achievement**, awarding entity or agency, (awarded amount, duration)

Service to the community & memberships in academic organization

- **Refereeing activity for scientific journals**, give concise details
- **Membership in relevant professional societies, consortia or collaborations**, give concise details
- **Function in associations**, give concise details

Extended scientific or exchange stays

- **Institution, program / purpose**, give concise details

Teaching Portfolio

List here the courses you have assisted in or taught until now.

- **Name of Course**, (type of course: e.g. lecture, exercises, practicum, ...) Semester 202X, ECTS of the course, Bachelor's / Master's Curriculum, list your co- or lead-lecturer, description of your responsibilities in this course.
- **Name of Course**, (type of course: e.g. lecture, exercises, practicum, ...) Semester 202X, ECTS of the course, Bachelor's / Master's Curriculum, list your co- or lead-lecturer, description of your responsibilities in this course.

Nota bene: If you use this template for your Habilitationsschrift, replace this section by the structure and content of the teaching portfolio, as recommended¹ by the *Zentrum für Lehrkompetenz (ZLK) of Graz University*.

Observing Portfolio

The candidate has obtained XXX nights of accumulated observing experience of at international facilities:

- Observatory name and location, telescope, technique, number of nights

Proposals lead as Principle investigator

- **Title of successful proposal**, awarded observing time (observing semester), observatory, telescope & instrument specification

Proposals contributed to as Co-investigator

- **Title of successful proposal**, awarded observing time (observing semester), name of PI, observatory, telescope & instrument specification

¹ <https://lehrkompetenz.uni-graz.at/de/service/lehrportfolios/>

List of Publications

Bibliographic information and overview

Repository identifiers	ORCID: xxx, GitHub: xxx
Bibliometric indicators	H-index: xxx, m-index: xxx
Publication record in NASA ADS	URL of your private NASA ADS library

Refereed journal publications

Published work that is available as open access through the ArXiv.org preprint server or journal policies is indicated through the abbreviation [AO].

1. **Your, N.**, Author, A., Author, B., Author, C., ... , **Title of the paper**, Journal, Volume, pages, DOI, (Year) [AO]
2. Author, A., Author, B., **Your, N.**, Author, C., ... , **Title of the paper**, Journal, Volume, pages, DOI, (Year) [AO]

Conference proceedings and other publications

1. **Your, N.**, Author, A., Author, B., Author, C., ... , **Title of the conference proceedings paper**, Conference Series, pages, DOI, (Year) [AO]
2. Author, A., Author, B., **Your, N.**, Author, C., ... , **Title of the popular paper**, Popular Journal, Volume, pages, DOI, (Year) [AO]

Colophon

This thesis was typeset with $\text{\LaTeX}$ 2 ϵ . It uses the *Clean Thesis* style developed by Ricardo Langner, available at <http://cleanthesis.der-ric.de/>. The design of the *Clean Thesis* style is inspired by user guide documents from Apple Inc.

The template is provided in this adapted version by Desmond Grossmann, BSc, Stefan Janisch, BSc, Lea Schimak and Paul Beck, PhD to implement the editorial guidelines of *Astronomy & Astrophysics* and the cooperate-identity guidelines of the University of Graz for the Institute for Physics. Version: March, 2024.

Declaration

I declare that I have authored this thesis independently, that I have not used other than the declared sources/resources, and that I have explicitly indicated all material which has been quoted either literally or by content from the sources used. The text document uploaded to UNIGRAZonline or TUGRAZonline is identical to the present master's thesis.

Graz, April 1, 2025

Markus Ornik